

Natpara®

(parathyroid hormone)
for Injection

25 • 50 • 75 • 100 mcg per dose strength



How you should store your NATPARA

What is NATPARA® (parathyroid hormone) for Injection?

- NATPARA is a prescription parathyroid hormone used with calcium and vitamin D to control low blood calcium (hypocalcemia) in people with low parathyroid hormone blood levels (hypoparathyroidism).
- NATPARA is only for people who do not respond well to treatment with calcium and active forms of vitamin D alone, because it may increase the possible risk of bone cancer (osteosarcoma).
- NATPARA was not studied in people with hypoparathyroidism caused by calcium-sensing receptor mutations.
- NATPARA was not studied in people who get sudden hypoparathyroidism after surgery.
- It is not known if NATPARA is safe and effective for children 18 years of age and younger. NATPARA should not be used in children and young adults whose bones are still growing.

Important Safety Information

What is the most important information I should know about NATPARA?

Warning: Possible bone cancer (osteosarcoma).

- During animal drug testing, NATPARA caused some rats to develop a bone cancer called osteosarcoma. It is not known if people who take NATPARA will have a higher chance of getting osteosarcoma. Tell your doctor right away if you have pain in any areas of your body that does not go away, or any new or unusual lumps or swelling under your skin that is tender to touch.
- **NATPARA is only available through the NATPARA Risk Evaluation and Mitigation Strategy (REMS) Program.** The purpose of the NATPARA REMS Program is to inform patients about the potential risk of osteosarcoma associated with the use of NATPARA. For more information about this REMS program, call 1-855-NATPARA (628-7272) or go to www.NATPARAREMS.com.

Please see additional Important Safety Information on the next page.

Please see full [Prescribing Information](#), [Medication Guide](#), and [Instructions for Use](#) for NATPARA.

How you should store your NATPARA

How should I store NATPARA?

- **Unmixed NATPARA medicine cartridges**
 - Refrigerate NATPARA between 36°F to 46°F (2°C to 8°C)
 - **Do not freeze**
- **Mixed NATPARA medicine cartridges**
 - Refrigerate NATPARA between 36°F to 46°F (2°C to 8°C)
 - **Do not freeze**
 - You can use the Q-Cliq™ pen for up to 14 days after mixing the medicine cartridge
 - Throw away the mixed NATPARA medicine cartridges 14 days after mixing the medicine cartridge
- Store NATPARA away from heat and light
- **Do not** freeze or shake NATPARA. **Do not** use NATPARA if it was frozen or shaken



Keep NATPARA and all medicines out of the reach of children.

Please see [Medication Guide](#) and [Instructions for Use](#) for information on how to use NATPARA.



Speak to your doctor or visit www.NATPARA.com to learn more.

Important Safety Information (cont'd)

What is the most important information I should know about NATPARA? (cont'd)

NATPARA may cause other serious side effects, including: (cont'd)

High blood calcium (hypercalcemia)

- NATPARA can cause some people to have a higher blood calcium level than normal.
 - Your doctor should check your blood calcium before you start and during your treatment with NATPARA.
 - Tell your doctor if you have nausea, vomiting, constipation, low energy, or muscle weakness. These may be signs that you have too much calcium in your blood.

Tell your doctor right away if you have any of these signs and symptoms of **high or low blood calcium** levels.

Low blood calcium (hypocalcemia)

- People who stop using or miss a dose of NATPARA may have an increased risk of severe low blood calcium levels.
- Tell your doctor if you have tingling of your lips, tongue, fingers and feet, twitching of face muscles, cramping of feet and hands, seizures, depression, or have problems thinking or remembering.

Who should not use NATPARA?

- **Do not use NATPARA** if you are allergic to parathyroid hormone or any of the ingredients in NATPARA.

What should I tell my healthcare provider before using NATPARA?

- **Before you start using NATPARA, tell your doctor about all of your medical conditions. Tell your doctor about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of NATPARA?

- NATPARA may cause serious side effects like allergic (hypersensitivity) reaction, including anaphylaxis. Tell your healthcare provider or get emergency medical help right away if you have any of the following symptoms of an allergic reaction:
 - swelling of your face, lips, mouth, or tongue
 - breathing problems
 - fainting, dizziness, feeling lightheaded (low blood pressure)
 - fast heartbeat
 - itching
 - rash
 - hives
- **The most common side effects of NATPARA include:** tingling, tickling, or burning feeling of the skin, low or high blood calcium, headache, nausea, reduced sense of touch or sensation, diarrhea, vomiting, pain in joints, too much calcium in urine, and pain in limbs.

These are not all the possible side effects of NATPARA. For more information, talk with your doctor. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit: www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full [Prescribing Information](#), [Medication Guide](#), and [Instructions for Use](#) for NATPARA.



©2024 Takeda Pharmaceuticals U.S.A., Inc., 500 Kendall Street, Cambridge, MA 02142. 1-877-TAKEDA-7 (1-877-825-3327). All rights reserved. TAKEDA® and the TAKEDA Logo® are registered trademarks of Takeda Pharmaceutical Company Limited. NATPARA® and the NATPARA Logo® are registered trademarks of Takeda Pharmaceutical Company Limited. Q-Cliq™ is a trademark of Becton-Dickinson. US-NAT-0280v2.0 03/24

 **Natpara**®
(parathyroid hormone)
for Injection
25 • 50 • 75 • 100 mcg per dose strength